

University of Groningen

Osteoprotegerin in Fibrotic Disorders

Adhyatmika, A.

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Adhyatmika, A. (2018). *Osteoprotegerin in Fibrotic Disorders*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter I

GENERAL INTRODUCTION

AIM AND SCOPE OF THE THESIS

Clinical studies have reported higher serum level of OPG in cases of fibrosis, especially liver fibrosis¹⁻³, but there is limited data available to explain the significance of this protein in the biology of the disease. This thesis is aimed to give wider insight into this topic.

In **chapter 2**, we studied the possible roles OPG could have in liver fibrosis using human samples, mouse models of liver fibrosis, murine precision-cut liver slices, and relevant cell lines. These studies included upregulation of its production by TGF β and investigation into its role as a decoy receptor of RANKL. In addition, we found that OPG itself can upregulate the expression of TGF β , which suggests that OPG and TGF β are involved in a feed-forward loop. In **chapter 3** we studied whether the role of OPG in liver fibrosis could be extended to a more general role in fibrotic processes. We investigated whether induction of fibrosis in other organs (lung, kidney, colon) also resulted in increased OPG production using precision-cut slices of these organs. Furthermore, we studied how OPG responds to antifibrotic treatment in the different organ slices. In **chapter 4** we further studied regulation of OPG by different cytokines and found that in addition to TGF β , stimulation with IL13 could also induce expression of OPG. We further investigated how IL13 can induce OPG production and found this is also mediated through TGF β . Hypothesizing a feedback mechanism controlling the TGF β -OPG feed-forward loop, in **chapter 5**, we investigated the influence of miR-145-5p in controlling the OPG-TGF β profibrotic loop.

OPG has an important role in the regulation of bone formation and degradation by inhibiting osteoclast activation and function in bone extracellular matrix degradation. We hypothesized a similar role for OPG in fibrosis by inhibiting antifibrotic macrophages and we therefore in **chapter 6** investigated literature to see what is known about antifibrotic macrophages and how we could look at those in the future studies involving OPG.

Finally, we discuss and summarize our findings in **chapter 7** with future perspectives of the potential use of OPG as a new therapeutic target or biomarker for (liver) fibrosis.

FIBROSIS

Fibrosis is abnormal fibrogenesis characterized by the accumulation of extracellular matrix molecules such as collagens and fibronectin produced by myofibroblasts that are characterized by increased expression of α -smooth muscle actin (α SMA)^{4,5}. The pathology of fibrosis is marked by the unbalanced regulation of inflammation, matrix formation, and resolution, resulting in maintaining persistent formation of non-functional matrix with no or very slow resolution towards functional tissue replacement^{6,7}. The disbalance can be caused by several factors, which can be related to each other: (1) continuous signalling from the injury or infection that trigger immune responses⁸, (2) irregular cellular functions or differentiation, for examples of macrophages and fibroblasts^{9,10}, and (3) homeostatic abnormalities caused by several possible factors such as a genetic disorder^{11,12} or the absence of blood vessels at the site of fibrosis¹³.

In most cases, fibrosis is associated with organ failure as its advanced pathological event¹⁴, and organ transplantation is then the only option for treatment¹⁵. Much effort has been put in by researchers in understanding the biology and cause of organ fibrosis in order to find a strategy to prevent and to cure. However, to date no effective treatment has been found yet¹⁶. In some specific types of fibrosis such as cystic fibrosis and familial pulmonary fibrosis, heredity is the cause of the disease^{17,18}. However, in most other cases of fibrosis, the disease develops over a lifetime and the exact cause cannot be determined anymore^{19,20}. Although there are some risk factors like chronic infection and chemical exposures that are known to increase the occurrence rate, it has not been proven yet whether they are the main cause of the disease.

Fibrosis can occur in many organs, including vital organs like the liver, the lungs, the heart, the intestines and the bone marrow. Of those organs, pulmonary and liver fibrosis are most prevalent^{21,22,23} and researchers and clinicians are trying to find a way to detect the disease in the earliest phase possible in order to prevent or to avoid an incurable stage of the disease.

LIVER FIBROSIS

Liver fibrosis is a burden to mankind, contributing to almost 2% of global deaths in 2010²⁴. The pathology of the disease is characterized by the accumulation of extracellular matrix consisting of collagens and other matrix proteins in the liver, preventing regeneration of new functional parenchymal cells²⁵. Liver fibrosis is thought to be the result of unbalanced wound healing process caused by several chronic factors such as hepatitis infection, alcohol abuse, or long-term use of hepatotoxic drugs (figure 1)²⁶.




Figure 1. Schematic of liver fibrosis progression and resolution as summarized by Pellicoro et al. (2014). The disease is hypothesized to start with chronic inflammation followed by loss of function and can ultimately end in cirrhosis²⁶.

Fibrotic sites in the liver are dominated by activated hepatic stellate cells instead of hepatocytes and these stellate cells produce profibrotic cytokines and growth factors to maintain the state of fibrosis²⁵. The early state of liver fibrosis is hard to diagnose since symptoms of disturbed liver function only occur when the liver is moderately to severely compromised²⁷. Although it is considered possible to reverse the fibrotic process towards resolution²⁸, to date there is no effective medical treatment available that can do this, especially in cases of full-blown cirrhosis²⁹. Therefore, liver transplantation is currently the only option to restore organ functionality in time.

The pathogenesis of liver fibrosis includes, but is not limited to, three main factors: (1) persistent inflammatory and immune responses by macrophages and lymphocytes from long-term injury³⁰, (2) hepatocyte death either by apoptosis or necrosis and replacement by fibroblasts³¹, and (3) high oxidative stress that triggers stellate cells activation, this is especially the case for fibrosis caused by infections and chemical injury³². The process of liver fibrosis is generally thought to start with continuous inflammation that activates Kupffer cells, liver resident macrophages, to release various proinflammatory and profibrotic cytokines such as TNF α , IL1 β , TGF β , and PDGFBB³³. This is followed by activation of hepatic stellate cells and transformation of these cells into myofibroblasts that are the main producers of the excess extracellular matrix that characterizes fibrosis³⁴. The activation of hepatic stellate cells is prominent and therefore is the most studied event for drug development purposes^{35,36}.

LIVER FIBROSIS RESOLUTION

It is a matter of debate among scientists whether the process of liver fibrosis can be reversed, especially when the disease has already developed. Current liver fibrosis medication is aimed at inhibiting inflammation to stop oxidative stress and other inflammatory stimulants and is thereby expected to consequently protect the remaining functional hepatocytes. Resolution is then expected to occur spontaneously. As inflammation is thought to be important in the process of fibrosis, the use of anti-inflammatory drugs, mainly corticosteroids, is widely proposed^{37,38}. However, this approach appears not to be effective and casts doubts on the importance of inflammation in the later stages of fibrosis³⁹. Recently, there are some other approaches under development, including degrading or altering the composition of extracellular matrix⁴⁰, inhibiting further hepatic stellate cells (HSCs) activation⁴¹, supporting hepatocytes growth using hepatocyte growth factor (HGF)⁴², and directing macrophages differentiation towards antifibrotic phenotypes⁴³. With these approaches, some experimental studies showed promising results towards resolution of liver fibrosis, marked by the lower expression of some frontline markers like collagen and α SMA⁴⁴.

As liver fibrosis is often represented as the activity of HSCs, the progression as well as resolution of liver fibrosis can be followed by simply evaluating the activation and deactivation or apoptosis of HSCs⁴¹. Both directions involve communication with lymphocytes and macrophages, mediated by several prominent mediators such as IFN γ , TGF β , IL13, and MMP9. Aforementioned markers such as collagen-1 and α SMA can be used to evaluate changes in HSCs activation and deactivation/apoptosis (figure 2)²⁶.




Figure 2. Liver fibrosis progression and resolution can be represented by HSCs activation and reversion. The process involves T cells and macrophages, communicating via various cytokines and growth factors with HSCs, and the process can be evaluated using several markers expressed by HSCs. The figure is republished with permission²⁶.

The proposed approaches to stimulate liver fibrosis resolution are generally to support antifibrotic machineries and to suppress profibrotic ones. In order to interfere within a persistent fibrotic condition and to induce resolution, knowledge of how the involved cells communicate with each other during normal resolution is essential. Targeting the key mediators involved in this cellular communication during fibrosis may solve the question how resolution can be brought about by medical effort. In order to reach that goal, better

understanding of the roles and activities of the various communication mechanisms is important.

Since liver fibrosis is a chronic process involving immune responses, inflammation, and wound healing, many soluble mediators have been reported to take part in the biology of the disease. The mediators can be divided into functional subgroups: Chemokines such as CCL2, CCL3, CCL17, CCL22, CXCL9, and CXCL10 that are responsible for cell recruitment to the site of fibrosis⁴⁵⁻⁴⁷; immune and inflammatory cytokines such as IL1 β , TNF α , IFN γ , and IL33 that are responsible for oxidative stress as well as macrophage and lymphocyte activation⁴⁸; and growth factors, cytokines and enzymes involved in tissue remodelling such as TGF β , PDGF, IL4, IL13, fibronectin, tissue transglutaminase, MMP9, and MMP13 that are responsible for the activation of fibroblasts, tissue regeneration, and resolution of fibrosis⁴⁹⁻⁵⁴. There are some proteins or compounds with the increased expression or production during liver fibrosis but not considered as mediators, because these are components of the extracellular matrix itself i.e. collagens, glycoproteins, and hyaluronic acid, which are also then considered as biomarkers⁵⁵. TGF β has been reported to be responsible for many profibrotic signals especially in liver fibrosis albeit it also has essential functions in homeostasis⁵⁶. TGF β signalling through the SMAD pathway activates hepatic stellate cells and contributes to lipid accumulation, therefore supports inflammation and maintains extracellular matrix⁵⁴. Moreover, TGF β also contributes to hepatocyte cell death and increasing ROS production to induce oxidative stress⁵⁷.

Recent reports have also shown that the bone matrix-related protein osteoprotegerin (OPG) in blood is increased during liver fibrosis and can serve as a biomarker to increase diagnostic accuracy of tests to diagnose liver fibrosis¹⁻³. Although involved in bone matrix regulation, OPG itself is not an extracellular matrix component and its role in liver fibrosis has not been elucidated. Interestingly, a recent study has reported that TGF β can induce OPG production in synovial fibroblasts in arthritis⁵⁸. This information suggests a link between OPG and fibroblasts and thereby suggests a potential role in the pathogenesis of liver fibrosis.

OSTEOPROTEGERIN IN FIBROSIS

Osteoprotegerin (OPG), also known as tumor necrosis factor receptor (TNF) superfamily 11B (TNFRS11B), has been widely studied as a decoy receptor of RANKL, also known as TNF ligand superfamily 11 (TNFSF11), and TRAIL, also known as TNF ligand superfamily 10 (TNFSF10)⁵⁹. OPG was first discovered to be produced by osteoblasts and regulates bone formation and remodelling by binding to RANKL thus preventing osteoclast activation⁶⁰. Osteoprotegerin is produced as a 60 kDa-monomer consisting 401 amino acids. The monomers may also be assembled at the cys-400 residue to form 120 kDa disulphide-linked dimers. Both monomer and dimer proteins have a signal peptide, which is cleaved prior to secretion to form active OPG. As a decoy receptor for RANKL and TRAIL, the structure of OPG consists of four cysteine-rich pseudo repeats located in the N-terminal, responsible for its binding activity to RANKL and TRAIL. However, it lacks a trans-membrane domain for attachment to cell membranes and is thus biologically available as a soluble protein, which increases its effectiveness to catch RANKL and TRAIL available in the environment^{61,62}.

Further studies have shown that OPG is not only produced by osteoblasts, but also smooth muscle cells, fibroblasts and cancer cells and is postulated to have a significant role in arthritis and cancer^{59,60,63}. In cancer, for instance, OPG is produced by cancer cells to intercept TRAIL to avoid TRAIL-receptor mediated apoptosis⁶⁴, while in arthritis OPG was shown to be produced by synovial-like fibroblasts to maintain their activated state as well as to avoid apoptosis⁶⁵.

Recent studies have also reported OPG to be higher in several types of organ fibrosis. Garcia-Valdecasas-Campelo (2006) reported higher serum OPG levels in patients with liver fibrosis and alcoholic liver disease¹, Boorsma *et al.* (2015) in patients with pulmonary fibrosis⁶⁶, and Sen *et al.* (2005) in patients with postoperative epidural fibrosis⁶⁷. Including OPG as an additional biomarker to a panel of biomarkers to diagnose liver fibrosis has been introduced by Bosselut *et al.* (2013) and proven to increase the diagnostic accuracy of their panel³. However, it is not clear what role OPG plays in fibrotic processes in general and liver fibrosis in particular.

In addition to the hypothesis of preventing TRAIL-induced apoptosis of activated myofibroblasts, another possible hypothesis is that OPG can interfere with break-down of extracellular matrix. Meng *et al.* reported that OPG directly inhibits the production of matrix metalloproteinase-13 (MMP13)⁶⁸, an important

antifibrotic MMP in liver fibrosis⁶⁹. Moreover, Corisdeo *et al.* (2001) suggested in their study that RANKL stimulates the production of cathepsin K, a collagen-degrading protease, in bone marrow cultures and macrophages⁷⁰. This finding suggests that high levels of OPG can prevent expression of MMP13 and cathepsin K in other cells like macrophages and thus inhibit ECM degradation. Furthermore, Toffoli *et al.* (2011) showed that OPG could promote vascular fibrosis by inducing TGF β 1 production in vitro and in vivo⁷¹.

However, despite the evidence of the possible profibrotic activities of OPG, there are no studies explaining how OPG contributes to the development of organ fibrosis especially in the liver. As a soluble receptor, there is very limited evidence of OPG directly interacting with a membrane receptor, therefore any profibrotic activity of OPG is mostly like explained by its abilities to scavenge ligands like RANKL and TRAIL.

MICRORNA AND THEIR ROLES IN FIBROSIS

MicroRNAs (miRNAs) are non-coding small RNA molecules that can degrade or block translation of their target messenger RNA (mRNA) sequences. This is done in collaboration with the protein Argonaut in a complex called RNA-induced silencing complex (RISC)⁷². miRNAs were first discovered by Lee *et al.* (1993) and Wightman *et al.* (1993) when studying negative regulation of lin-14 in *Caenorhabditis elegans* through formation of small molecule RNA later described as microRNA^{73,74}. From then on, miRNAs have been studied widely as feedback mechanisms of many biological mechanisms in various diseases especially in cancer and fibrosis⁷⁵. Particularly during fibrosis, miRNAs can be either up- or down-regulated, and thereby contribute to the pathogenesis, progression, and resolution of fibrosis (*figure 3*)^{76,77}.

The role of OPG in liver fibrosis is very likely to also involve microRNAs in some way. Ong *et al.* have recently reported microRNA expression levels in human fibroblasts with or without TGF β treatment, and found that some of the TGF β -upregulated miRNAs can target OPG, one of those being miR145-5p⁷⁸.




Figure 3. The up- and down-regulation of miRNAs in idiopathic pulmonary fibrosis (IPF) as summarized by Li et al. (2016). As miRNAs can block cytokine expression, miRNAs can play pro- and antifibrotic roles in fibrosis. MiRNAs can interfere with the activities of many cells like epithelial cells, macrophages, and fibroblasts. Changes in the expression of miRNAs during fibrosis can alter homeostasis and therefore, these molecules may be potential targets for therapy. Figure is reprinted with permission⁷⁷.

At least three different recent studies have reported effects of miR-145 on expression of OPG. Jia et al. (2017) reported that transfecting human osteoblast-like MG-63 cells with miR-145-5p mimics can partially inhibit OPG upregulation by estrogen⁷⁹. In another study, Wang et al. (2017) confirmed OPG as a direct target for mir-145 by using a dual-luciferase reporter assay and further showed that higher mir-145 expression can suppress OPG expression⁸⁰. Finally, Zhao et al. (2016) showed that OPG expression was significantly lower after lentivirus-mediated transfection of rats and THP-1 cells with miR-145⁸¹.

As miR-145 obviously targets OPG, it may also be involved in regulation of OPG production in fibroblasts. Interestingly, Yang et al. (2013) reported upregulation of miR-145 in TGFβ1-treated lung fibroblasts as compared to untreated cells⁸². Furthermore, Zhou et al. (2016) reported in their study that miR-145 inhibited TGFβ-induced human and rat hepatic stellate cells activation and proliferation in vitro and downregulation of this microRNA in vivo in CCl₄-induced liver fibrosis in mice⁸³.

Based on literature reports, we hypothesized that OPG is produced by fibroblasts in liver tissue and that OPG is associated with fibrotic processes because its expression is controlled by TGF β . In addition, this expression may be further regulated by microRNAs, especially miR-145.

REFERENCES

1. García-Valdecasas-Campelo E, González-Reimers E, Santolaria-Fernández F, De La Vega-Prieto MJ, Milena-Abril A, Sanchez-Perez MJ, Martínez-Riera A, and Gómez-Rodríguez MDLA, 2006, Serum osteoprotegerin and RANKL levels in chronic alcoholic liver disease, *ALCOHOL & ALCOHOLISM*, **41**(3): 261-66
2. Yilmaz Y, Yonal O, Kurt R, Oral AY, Eren F, Ozdogan O, Ari F, Celikel CA, Korkmaz S, Ulukaya E, Imeryuz N, Kalayci C, and Avsar E, 2010, Serum levels of osteoprotegerin in the spectrum of nonalcoholic fatty liver disease, *Scandinav J Clin Lab Invest*, **70**: 541-46
3. Bosselut N, Taibi L, Guéchet J, Zarski JP, Sturm N, Gelineau MC, Poggi B, Thoret S, Lasnier E, Baudin B, Housset C, Vaubourdolle M, and the ANRS HCEP 23 Fibrostar Group, 2013, Including osteoprotegerin and collagen IV in a score-based blood test for liver fibrosis increases diagnostic accuracy, *Clin Chim Acta*, **415**: 63-8
4. Ho YY, Lagares D, Tager AM, and Kapoor M, 2014, Fibrosis—a lethal component of systemic sclerosis, *Nat Rev Rheum*, **10**: 390-402
5. Boukhalfa G, Desmoulière A, Rondeau E, Gabbiani G, and Sraer JD, 1996, Relationship between alpha-smooth muscle actin expression and fibrotic changes in human kidney, *Eur PMC*, **4**(4): 241-47
6. Lee SB and Kalluri R, 2010, Mechanistic connection between inflammation and fibrosis, *Kidney Intl Suppl*, **119**: S22-S26
7. Wynn TA, 2008, Cellular and molecular mechanisms of fibrosis, *J Pathol*, **214**(2): 199-210
8. Wick G, Backovic A, Rabensteiner E, Plank N, Schwendtner C, and Sgonc R, 2010, The immunology of fibrosis: innate and adaptive responses, *Trends Immunol*, **31**(3): 110-9
9. Tacke F and Zimmermann HW, 2014, Macrophage heterogeneity in liver injury and fibrosis, *J Hepatol*, **60**(5): 1090-6
10. Darby IA and Hewitson TD, 2007, Fibroblast differentiation in wound healing and fibrosis, *Int Rev Cytol*, **257**: 143-79
11. Cutting GR, 2015, Cystic fibrosis genetics: from molecular understanding to clinical application, *Nat Rev Gen*, **16**: 45-56
12. Evans CM, Fingerlin TE, Schwarz MI, Lynch D, Kurche J, Warg L, Yang IV, and Schwartz DA, 2016, Idiopathic pulmonary fibrosis: A genetic disease that involves mucociliary dysfunction of the peripheral airways, *Physiol Rev*, **96**(4): 1567-91
13. Mann CJ, Perdiguer E, Kharraz Y, Aguilar S, Pessina P, Serrano AL, and Muñoz-Cánoves P, 2011, Aberrant repair and fibrosis development in skeletal muscle, *Skelet Muscle*, **1**:21
14. Rockey DC, Bell PD, and Hill JA, 2015, Fibrosis—A common pathway to organ injury and failure, *New Engl J Med*, **372**: 1138-49
15. Said A and Lucey MR, 2008, Liver transplantation: an update 2008, *Curr Opin Gastroenterol*, **24**: 339-45
16. Rosenbloom J, Mendoza FA, and Jimenez SA, 2013, Strategies for anti-fibrotic therapies, *Biochim Biophys Acta*, **1832**(7): 1088-103
17. Riordan JR, Rommens JM, Kerem B, Alon N, Rozmahel R, Grzelczak Z, Zielenski J, Lok S, Plavsic N, Chou JL, et al., 1989, Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA, *Science*, **245**(4922): 1066-73
18. Garcia CK, 2011, Idiopathic pulmonary fibrosis: update on genetic discoveries, *Proc Am Thorac Soc*, **8**(2): 158-62
19. Bettler and Brenner, 2005, Liver fibrosis, *J Clin Invest*, **115**(2): 109-218
20. Gross TJ and Hunninghake GW, 2001, Idiopathic pulmonary fibrosis, *New Engl J Med*, **345**: 517-25
21. Blachier M, Leleu H, Peck-Radosavljevic M, Valla DC, and Roudot-Thoraval F, 2013, The burden of liver disease in Europe: A review of available epidemiological data, *J Hepatol*, **58**(3): 593-608
22. Göbel T, Erhardt A, Herwig M, Poremba C, Baldus SE, Sagir A, Heinzel-Pleines U, and Häussinger D, 2011, High prevalence of significant liver fibrosis and cirrhosis in chronic hepatitis B patients with normal ALT in central Europe, *J Med Virol*, **83**(6): 968-73
23. Nalysnyk L, Cid-Ruzafa J, Rotelia P, and Esser D, 2012, Incidence and prevalence of idiopathic pulmonary fibrosis: review of the literature, *Eur Respir Rev*, **21**: 355-61
24. Mokdad AA, Lopez AD, Shahrzaz S, Lozano R, Mokdad AH, Stanaway J, Murray CJL, and Naghavi M, 2014, Liver cirrhosis mortality in 187 countries between 1980 and 2010: a systemic analysis, *BMC Med*, **12**:145
25. Hernandez-Gea V and Friedman SL, 2011, Pathogenesis of liver fibrosis, *Annu Rev Pathol Mech Dis*, **6**: 425-56
26. Pellicoro A, Ramachandran P, Iredale JP, and Fallowfield JA, 2014, Liver fibrosis and repair: immune regulation of wound healing in a solid organ, *Nat Rev Immunol*, **14**: 181-94

27. Tsochatzis EA, Bosch J, and Burroughs AK, 2014, Liver cirrhosis, *Lancet*, **383**(9930): 1749-61
28. Ismail MH and Pinzani M, 2009, Reversal of liver fibrosis, *Saudi J Gastroenterol*, **15**(1): 72-9
29. Ge PS and Runyon BA, 2016, Treatment of patients with cirrhosis, *N Engl J Med*, **375**: 767-77
30. Robinson MW, Harmon C, and O'Farrelly C, 2016, Liver immunology and its role in inflammation and homeostasis, *Cell Mol Immunol*, **13**(3): 267-76
31. Guicciardi ME, Malhi H, Mott JL, and Gores GJ, 2013, Apoptosis and necrosis in the liver, *Compr Physiol*, **3**(2): 977-1010
32. Lee KS, Lee SJ, Park HJ, Chung JP, Han KH, Chon CY, Lee SI, and Moon YM, 2001, Oxidative stress effect on the activation of hepatic stellate cells, *Yonsei Med J*, **42**(1): 1-8
33. Ramachandran P and Iredale JP, 2012, Macrophages: central regulators of hepatic fibrogenesis and fibrosis resolution, *J Hepatol*, **56**(6): 1417-9
34. Lee UE and Friedman SL, 2011, Mechanisms of hepatic fibrogenesis, *Best Pract Res Clin Gastroenterol*, **25**(2): 195-206
35. Wu J and Zern MA, 2000, Hepatic stellate cells: a target for the treatment of liver fibrosis, *J Gastroenterol*, **35**(9): 665-72
36. Bataller R and Brenner DA, 2001, Hepatic stellate cells as a target for the treatment of liver fibrosis, *Semin Liver Dis*, **21**(3): 437-51
37. Reyes-Gordillo K, Shah R, and Muriel P, 2017, Oxidative stress and inflammation in hepatic diseases: Current and future therapy, *Oxid Med Cell Longev*, doi: 10.1155/2017/3140673
38. Czaja AJ, 2014, Hepatic inflammation and progressive liver fibrosis in chronic liver disease, *World J Gastroenterol*, **20**(10): 2515-32
39. Trivedi PJ and Hirschfield GM, 2013, Treatment of autoimmune liver disease: current and future therapeutic options, *Ther Adv Chronic Dis*, **4**(3): 119-41
40. Klaas M, Kangur T, Viil J, Mäemets-Allas K, Minajeva A, Vadi K, Antsov M, Lapidus N, Järvekülg, and Jaks V, 2016, The alterations in the extracellular matrix composition guide the repair of damaged liver tissue, *Sci Rep*, **6**: 27398
41. Zhang CY, Yuan WG, He P, Lei JH, and Wang CX, Liver fibrosis and hepatic stellate cells: ethiology, pathological hallmarks, and therapeutic targets, *World J Gastroenterol*, **22**(48): 10512-22
42. Xia JL, Dai C, Michalopoulos GK, and Liu Y, 2006, Hepatocyte Growth Factor attenuates liver fibrosis induced by bile duct ligation, *Am J Pathol*, **168**(5): 1500-12
43. Adhyatmika A, Putri KSS, Beljaars L, and Melgert BN, 2015, The elusive antifibrotic macrophage, *Front Med (Lausanne)*, doi: 10.3389/fmed.2015.00081
44. Liu X, Xu J, Brenner DA, and Kisseleva T, 2013, Reversibility of liver fibrosis and inactivation of fibrogenic myofibroblasts, *Curr Pathobiol Rep*, **1**(3): 209-14
45. Ehling J, Bartneck M, Wei X, Gremse F, Fech V, Möckel D, Baeck C, Hittatiya K, Eulberg D, Luedde T, Kiessling F, Trautwein C, Lammers T, and Tacke F, 2014, CCL2-dependent infiltrating macrophages promote angiogenesis in progressive liver fibrosis, *Gut*, **63**(12): 1960-71
46. Heinrichs D, Berres ML, Nellen A, Fischer P, Scholten D, Trautwein C, Wasmuth HE, and Sahin H, The chemokines CCL3 promotes experimental liver fibrosis in mice, *PLoS ONE*, **8**(6): e66106
47. Zeremski M, Dimova R, Astemborski J, Thomas DL, and Talal AH, 2011, CXCL9 and CXCL10 chemokines as predictors of liver fibrosis in a cohort of primarily African-American injection drug users with chronic hepatitis C, *J Infect Dis*, **204**(6): 832-6
48. Borthwick LA, Wynn TA, and Fisher AJ, 2013, Cytokine mediated tissue fibrosis, *Biochim Biophys Acta*, **1832**(7): 1049-60
49. Hellerbrand C, Stefanovic B, Giordano F, Burchardt ER, and Brenner DA, 1999, the role of TGFbeta1 in initiating hepatic stellate cell activation in vivo, *J Hepatol*, **30**(1): 77-87
50. Kinnman N, Gorla O, Wendum D, Gendron MC, Rey C, Poupon R, and Housset C, 2001, Hepatic stellate cell proliferation is an early platelet-derived growth factor-mediated cellular event in rat cholestatic liver injury, *Lab Invest*, **81**(12): 1709-16
51. Sugimoto R, Enjoji M, Nakamuta M, Ohta S, Kohjima M, Fukushima M, Kuniyoshi M, Arimura E, Morizono S, Kotoh K, and Nawata H, 2005, Effect of IL-14 and IL-13 on collagen production in cultured human LI90 human hepatic stellate cells, *Liver Int*, **25**(2): 420-8
52. To WS and Midwood KS, 2011, Plasma and cellular fibronectin: distinct and independent functions during tissue repair, *Fibrogenesis Tissue Repair*, **4**:21
53. Haroon ZA, Hettasch JM, Lai TS, Dewhirst MW, and Greenberg CS, 1999, Tissue transglutaminase is expressed, active, and directly involved in rat dermal wound healing and angiogenesis, *FASEB J*, **13**(13): 1787-95
54. Caley MP, Martins VLC, and O'Toole EA, 2015, Metalloproteinases and wound healing, *Adv Wound Care (New Rochelle)*, **4**(4): 225-34
55. Fontana RJ, Goodman ZD, Dienstag JL, Bonkovsky HL, Naishadham D, Sterling RK, Su GL, Ghosh M, Wright EC, and the HALT-C Trial Group, 2008,

- Relationship of serum fibrosis markers with liver fibrosis stage and collagen content in patients with advanced chronic hepatitis C, *HEPATOLOGY*, **47**(3): 789-98
56. Meng XM, Nikolic-Paterson DJ, and Lan HY, 2016, TGF- β : the master regulator of fibrosis, *Nat Rev Nephrol*, **12**: 325-38
 57. Yang L, Roh YS, Song J, Zhang B, Liu C, Loomba R, and Seki E, 2014, Transforming growth factor beta signaling in hepatocytes participates in steatohepatitis through regulation of cell death and lipid metabolism in mice, *HEPATOLOGY*, **59**(2): 483-95
 58. Tunyogi-Csapo M, Kis-Toth K, Radacs M, Farkas B, Jacobs JJ, Finnegan A, Mikecz K, and Glant TT, 2008, Cytokine-controlled RANKL and osteoprotegerin expression by human and mouse synovial fibroblasts (Fibroblast-mediated pathologic bone resorption), *Arthritis & Rheumatism*, **58**(8): 2397-408
 59. Vitovski S, Phillips JS, Sayers J, and Croucher PJ, 2007, Investigating the interaction between osteoprotegerin and receptor activator of NF κ B or tumor necrosis factor-related apoptosis-inducing ligand: evidence for a pivotal role for osteoprotegerin in regulating two distinct pathways, *J Biol Chem*, **282**(43): 31601-609
 60. Boyce BF and Xing L, 2007, Biology of RANK, RANKL, and osteoprotegerin, *Arth Res Ther*, **9**(Suppl 1): S1
 61. Wright HL, McCharty HS, Middleton J, and Marshall MJ, 2009, RANK, RANKL, and osteoprotegerin in bone biology and disease, *Curr Rev Musculoskelet Med*, **2**(1): 56-64
 62. Anderson DM, Maraskovsky E, Billingsley WL, Dougall WC, Tometsko ME, Roux ER, Teepe MC, DuBose DC, Galibert L, 1997, A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function, *NATURE*, **390**:175-9
 63. Goswami S and Sharma-Walia N, 2015, Osteoprotegerin secreted by inflammatory and invasive breast cancer cells induces aneuploidy, cell proliferation and angiogenesis, *BMC Cancer*, **15**:935
 64. Lane D, Matte I, Rancourt C, and Piche A, 2012, Osteoprotegerin (OPG) protects ovarian cancer cells from TRAIL-induced apoptosis but does not contribute to malignant ascites-mediated attenuation of TRAIL-induced apoptosis, *J Ovarian Res*, **5**:34
 65. Miyashita T, Kawakami A, Nakashima T, Yamasaki S, Tamai M, Tanaka F, Kamachi M, Ida H, Migita K, Origuchi T, Nakao K, and Eguchi K, 2004, Osteoprotegerin (OPG) acts as an endogenous decoy receptor in tumour necrosis factor-related apoptosis-inducing ligand (TRAIL)-mediated apoptosis of fibroblast-like synovial cells, *Clin Exp Immunol*, **137**(2): 430-436
 66. Boersma CE, Draijer C, Cool R, Brandsma C, Nossent G, Brass DM, Times W, and Melgert BM, 2014, A possible role for the RANK/RANKL/OPG axis in pulmonary fibrosis, *Am J Respir Crit Care Med*, **189**: A1252
 67. Sen O, Gokcel A, Kizilkilic O, Erdogan B, Aydin MV, Sezgin N, Yalcin O, Caner H, and Altinors N, 2005, The relation between serum levels of osteoprotegerin and postoperative epidural fibrosis in patients who underwent surgery for lumbar disc herniation, *Neurol Res*, **27**(4): 452-455
 68. Meng H, Bai X, Yu H, Wang Z, and G Ai, 2017, Osteoprotegerin promotes cell growth by regulating matrix metalloproteinase-13 in chondrocytes, *J Biomat Tissue Eng*, **7**(3): 257-260
 69. Kim EJ, Cho HJ, Park D, Kim JY, Kim YB, Park TG, Shim CK, and Oh YK, 2011, Antifibrotic effect of MMP13-encoding plasmid DNA delivered using polyethylenimine shielded with hyaluronic acid, *Mol Ther*, **19**(2): 355-361
 70. Corisdeo S, Gyda M, Zaidi M, Moonga BS, and Troen BR, 2001, New insight into the regulation of cathepsin K gene expression by osteoprotegerin ligand, *Biochem Biophys Res Commun*, **285**(2): 335-39
 71. Toffoli B, Pickering RJ, Tsorotes D, Wang B, Bernardi S, Kantharidis P, Fabris B, Zauli G, Secchiero P, and Thomas MC, 2011, Osteoprotegerin promotes vascular fibrosis via a TGF- β 1 autocrine loop, *ATHEROSCLEROSIS*, **218**: 61-8
 72. Meister G, 2013, Argonaute proteins: functional insights and emerging roles, *Nat Rev Gen*, **14**: 447-459
 73. Lee RC, Feinbaum RL, and Ambros V, 1993, The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14, *CELL*, **75**(5): 843-54
 74. Wightman B, Ha I, and Ruvkun G, 1993, Posttranscriptional regulation of the heterochronic gene lin-14 by lin-4 mediates temporal pattern formation in C. elegans, *CELL*, **75**(5): 855-62
 75. Peng Y and Croce CM, 2016, The role of microRNAs in human cancer, *Sign Transduct Target Ther*, **1**:15004
 76. O'Reilly S, 2016, MicroRNAs in fibrosis: opportunities and challenges, *Arthritis Res Ther*, **18**:11
 77. Li H, Zhao X, Shan H, and Liang H, 2016, MicroRNAs in idiopathic pulmonary fibrosis: involvement in pathogenesis and potential use in diagnosis and therapeutics, *Acta Pharm Sin B*, **6**(6): 531-39
 78. Ong J, Timens W, Rajendran V, Algra A, Spira A, Lenburg ME, Campbell JD, van der Berge M,

- Postma DS, van den Berg A, Kluiver J, and Brandsma CA, 2017, Identification of transforming growth factor-beta-regulated microRNAs and the microRNA targetomes in primary lung fibroblasts, *PLOS One*, **doi**: 10.1371/journal.pone.0183815
79. Jia J, Zhou H, Zeng X, and Feng S, 2017, Estrogen stimulates osteoprotegerin expression via the suppression of mir-145 expression in MG-63 cells, *Mol Med Rep*, **15**(4): 1539-1546
80. Wang GD, Zhao XW, Zhang YG, Kong Y, Niu SS, Ma LF, and Zhang YM, 2017, Effects of mir-145 on the inhibition of chondrocyte proliferation and fibrosis by targeting TNFRSF11B in human osteoarthritis, *Mol Med Rep*, **15**(1): 75-80, **doi**: 10.3892/mmr.2016.5981
81. Zhao JJ, Wu ZF, Wang L, Feng DH, and Cheng L, 2016, MicroRNA-145 mediates steroid-induced necrosis of the femoral head by targeting the OPG/RANK/RANKL signalling pathway, *PLoS ONE*, **11**(7): e0159805
82. Yang S, Cui H, Xie N, Icyuz M, Banerjee S, Antony VB, Abraham E, Thannickal J, and Liu G, 2013, miR-145 regulates myofibroblast differentiation and lung fibrosis, *FASEB J*, **27**(6): 2382-2391
83. Zhou DD, Wang X, Wang Y, Xiang XJ, Liang ZC, Zhou Y, Xu A, Bi CH, and Zhang L, 2016, MicroRNA-145 inhibits hepatic stellate cell activation and proliferation by targeting ZEB2 through Wnt/ β -catenin pathway, *Mol Immunol*, **75**: 151-160